

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr MAVENCLAD®

Cladribine 10 mg tablets

Read this carefully before you start taking **MAVENCLAD** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **MAVENCLAD**.

What is MAVENCLAD used for?

- MAVENCLAD is used to treat adult patients with relapsing-remitting multiple sclerosis (RRMS). Multiple sclerosis (MS) is a disease that affects the brain and spinal cord.
- Treatment with MAVENCLAD has been shown to reduce the frequency of relapses and to slow down disability progression.
- MAVENCLAD is generally recommended for MS patients who have not responded well to, or cannot tolerate one or more of the other therapies for MS.

How does MAVENCLAD work?

Cladribine, the active substance in MAVENCLAD is cytotoxic, which means it causes cell death. Cladribine acts in specific ways on cells in your immune system called B and T lymphocytes. Cladribine helps to reduce inflammation in your body as a result of having MS.

What are the ingredients in MAVENCLAD?

Medicinal ingredients: cladribine

Non-medicinal ingredients: hydroxypropyl betadex, sorbitol and magnesium stearate. May contain trace amounts of sodium hydroxide and/or hydrochloric acid.

MAVENCLAD comes in the following dosage forms:

10 mg tablet

Do not use MAVENCLAD if:

- are allergic (hypersensitive) to cladribine or any of the other ingredients of MAVENCLAD;
- are at risk for infections because you have a weak immune system due to:
 - treatments you receive (for cancer [medication or radiation], chronic corticosteroids, bone marrow transplant(s), etc.);
 - a medical condition you have (for example, if you are infected with the human immunodeficiency virus [HIV]);
- have an active or inactive (past) infection, for example tuberculosis or liver inflammation (hepatitis);

- have or had a type of rare infection of the brain called progressive multifocal leukoencephalopathy (PML);
- have an active cancer;
- have moderate or severe kidney problems;
- are pregnant or breast-feeding;
- are a woman of childbearing potential or a man who could potentially father a child and you are not using birth control.

Talk to your healthcare professional if you are unsure if any of the above applies to you.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take MAVENCLAD. Talk about any health conditions or problems you may have, including if you:

- have a weakened immune system due to:
 - treatments you receive (for cancer [medication or radiation], bone marrow transplant(s), etc.);
 - a medical condition you have (for example, if you are infected with the human immunodeficiency virus [HIV]);
- have an infection;
- have liver problems;
- have recently been vaccinated;
- have or have previously had cancer;
- require blood transfusions;
- are pregnant, are thinking of becoming pregnant as MAVENCLAD may harm your baby;
- are breast-feeding or plan to breast-feed;
 - it is not known if MAVENCLAD goes into breast milk
 - do not breast-feed on the days you take MAVENCLAD, and for 10 days after the last dose
- are less than 18 years of age

Other warnings you should know about:

Blood Tests

You will have blood tests taken before you start treatment to see if you can take MAVENCLAD. Your healthcare professional will also do blood tests during and after treatment to ensure you can continue to take MAVENCLAD. The tests will determine if you are developing any complications from the treatment.

General Infections

You will be tested to see if you have any infections prior to starting MAVENCLAD. It is important to talk to your healthcare professional if you think you have an infection before, during or after treatment.

Symptoms of infections can include:

- fever
- aching, painful muscles
- headache
- generally feeling unwell
- loss of appetite

Your healthcare professional may delay start of treatment, or interrupt current treatment, until the infection clears up.

Vaccination

Talk to your healthcare professional if you have recently been given, or might be given, certain vaccines (called 'live' or 'live attenuated' vaccines).

- Treatment with MAVENCLAD can only begin 6 weeks after you have been vaccinated.
- If necessary, you will be vaccinated against varicella (chickenpox) prior to starting treatment.
- You must not be vaccinated during treatment with MAVENCLAD.
- You may only be vaccinated after MAVENCLAD treatment when your white blood cell count is normal.

Progressive Multifocal Leukoencephalopathy (PML)

Talk to your healthcare professional if you believe your MS is getting worse or if you notice any new symptoms, such as:

- weakness on one side of the body that gets worse
- clumsiness of your arms and legs
- trouble with your vision
- changes in your thinking and memory that lead to confusion and personality changes

These may be the symptoms of a rare brain disorder caused by infection and called progressive multifocal leukoencephalopathy (PML). PML is a serious condition that may lead to severe disability or death.

As a precaution, you may have a head MRI (magnetic resonance imaging) prior to starting treatment. This is particularly important if you have previously taken other multiple sclerosis treatments where PML is a risk.

Risk of Cancer

MAVENCLAD causes damage to the DNA (genotoxicity) and suppresses the immune system. As a result, MAVENCLAD could potentially increase the risk of cancer. In studies with MAVENCLAD, there were single cases of patients who developed different types of cancer, such as skin, ovarian, thyroid, breast and pancreatic cancer.

Follow your healthcare professional's instructions for screening for cancer.

Fructose Intolerance

MAVENCLAD contains sorbitol. Do not take MAVENCLAD if you have hereditary problems of fructose intolerance.

Pregnancy and Fertility

- It is important that:
 - women taking MAVENCLAD prevent pregnancy during treatment and for at least 6 months after the last dose.
 - men taking MAVENCLAD prevent pregnancy in their female partner while they are taking MAVENCLAD and for at least 6 months after the last dose.

- For female patients, it is not known if MAVENCLAD will reduce the efficacy of birth control pills. Given this, a barrier method of contraception (for example, condoms) should be added during treatment with MAVENCLAD and for at least 4 weeks after the last dose in each treatment year.
- If you or your partner gets pregnant during treatment or within the 6 months after the last dose, there may be a risk to the unborn baby. Your healthcare professional will stop treatment with MAVENCLAD if you are a woman and you get pregnant while taking it.
- MAVENCLAD may affect male fertility. Talk to your healthcare professional for more information.
- For male patients, it is not known if MAVENCLAD is detectable in semen. A barrier method of contraception (for example, condoms) should be used during treatment with MAVENCLAD and for 10 days after the last dose.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with MAVENCLAD:

Do not take MAVENCLAD together with medicines that weaken your immune system or your bone marrow. These include:

- cyclosporine, cyclophosphamide or azathioprine, used to suppress the immune system, for example after organ transplantation;
- methotrexate, used to treat conditions such as psoriasis or rheumatoid arthritis;
- long-term corticosteroids, used to reduce inflammation, for example in asthma. (Short-term corticosteroids can be used when advised by your healthcare professional).

Do not take MAVENCLAD at the same time as any other medicine. It may interact with other medicines in your stomach. Take other medicines 3 hours before or after you take MAVENCLAD.

Talk to your healthcare professional, if you are or have been treated with:

- medicines which may affect your blood cells (for example, carbamazepine used to treat convulsions). Your healthcare professional may need to supervise you more closely.
- dipyridamole, dilazep, nifedipine, nimodipine, reserpine, cilostazol or sulindac, used to treat the heart, high blood pressure, vascular conditions or inflammation
- cyclosporine (used to prevent organ transplant rejection)
- gefitinib, imatinib (used to treat cancer)
- elvitegravir, cobicistat (used to treat HIV/AIDS)
- eltrombopag (used to treat blood disorders)
- long-term corticosteroids (used to suppress inflammation)

How to take MAVENCLAD:

A. Understanding the MAVENCLAD Packaging

Follow the steps below for opening the MAVENCLAD package and for how to handle the tablets. As MAVENCLAD is cytotoxic, it is important that you follow the instructions below to make sure you handle and take MAVENCLAD safely.

1. Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).
2. Pick up the carton with the instructions facing up.
3. Open the package as follows (see Figure A below):
 - i. Open the flap on the left end.
 - ii. Push in both hooks on the sides of the carton. Use your index finger and thumb and keep the hooks pushed in.
 - iii. Pull the tray out until it stops. Do not try to remove the tray from the carton. It is important the tray stays in the carton to help prevent anyone from touching the tablets, especially children.




Figure A

4. Take the Package Insert out from the tray and read through all of it before taking MAVENCLAD.
5. Raise the blister pack by pushing your finger through the hole in the tray (see Figure B below). Place your hand under the blister pack and push 1 or 2 tablet(s) into your hand, based on your prescribed dose.




Figure B

6. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Swallow the tablet(s) with water immediately after removing them from the blister.
 - i. Do not leave your tablet(s) exposed on surfaces, for example on a table, or handle the tablet longer than necessary.
 - ii. If a tablet is left on a surface or if a broken or fragmented tablet is released from the blister, the area must be thoroughly washed.
 - iii. If you lose a tablet, contact your healthcare professional for advice.
7. **Wash your hands thoroughly with soap and water before touching your nose, eyes, and other parts of the body.**
8. Push the tray back into the carton.
9. Keep your tablets in the blister until your next dose. It is important that the tablets stay in the blister to help prevent anyone from touching the tablets or taking it by mistake.

B. Understanding Your Treatment Courses

You will be given MAVENCLAD as **two treatment courses over 2 years**.

One treatment course is **2 treatment weeks**, which are taken one month apart at the beginning of each treatment year.

A **treatment week** consists of **1 or 2 tablets daily given for 4 or 5 days**.

Example: if you start your treatment mid-April, you take your tablets as shown.

	Year 1		Year 2
1 st treatment week	1 or 2 tablets daily for 4 or 5 days, mid-April	1 st treatment week	1 or 2 tablets daily for 4 or 5 days, mid-April
2 nd treatment week)	1 or 2 tablets daily for 4 or 5 days, mid-May	2 nd treatment week	1 or 2 tablets daily for 4 or 5 days, mid-May

Before you start a treatment course, your healthcare professional will do a blood test to see whether the levels of lymphocytes (a type of white blood cells) are in an acceptable range. If this is not the case, your treatment will be delayed.

Once you have completed the 2 treatment years, your healthcare professional will observe your condition for another 2 years, in which you do not need to take the medicine. The efficacy of taking MAVENCLAD for longer than 2 years has not yet been established.

Usual dose:

Always take MAVENCLAD exactly as your healthcare professional has told you. Check with your healthcare professional if you are not sure. Take the tablet(s) at about the same time each day. You do not have to take the tablets at meal times. You can take them with or without food. Your healthcare professional may recommend that you take any other oral medication 3 hours before or after you take MAVENCLAD.

1. You will be prescribed the correct number of tablets for each treatment week, based on your body weight according to the table below.
2. In the left column of the table below find the row that fits your body weight (in kg), and then check the number of tablets that should be in the pack(s) for the treatment week you will be starting.
3. If the number of tablets in your pack(s) is different from the number shown for your weight in the table below, speak to your healthcare professional.
4. Note that for some weight ranges the number of tablets may vary from one treatment week to the next.

Example: if you weigh 85 kg and are about to start treatment week 1, you will be given 8 tablets.

Your weight in	Number of tablets to take			
	Year 1 treatment course		Year 2 treatment course	
kg	Treatment week 1	Treatment week 2	Treatment week 1	Treatment week 2
Less than 40 kg	Your healthcare professional will tell you the number of tablets to take			
40 to < 50	4	4	4	4
50 to < 60	5	5	5	5
60 to < 70	6	6	6	6
70 to < 80	7	7	7	7
80 to < 90	8	7	8	7
90 to < 100	9	8	9	8
100 to < 110	10	9	10	9
110 and above	10	10	10	10

Duration of a treatment week

Depending on the total number of tablets you have been prescribed, you have to take them over 4 or 5 days, in each treatment week.

The table below shows how many tablets (1 or 2 tablets) you have to take on each day. If your daily dose is 2 tablets, take them at the same time.

Example: if you have to take 8 tablets, you would take **2 tablets** on Day 1, Day 2, Day 3, then **1 tablet** on Day 4 and Day 5.

Total number of tablets per treatment week	Day 1	Day 2	Day 3	Day 4	Day 5
4	1	1	1	1	0
5	1	1	1	1	1
6	2	1	1	1	1
7	2	2	1	1	1
8	2	2	2	1	1
9	2	2	2	2	1
10	2	2	2	2	2

If you have any further questions on the use of this medicine, ask your healthcare professional.

Overdose:

There is limited experience with overdose of MAVENCLAD. It is known that the more of this medicine you take the fewer white blood cells you may have in your body.

If you think you, or a person you are caring for, have taken too much MAVENCLAD, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose and you remember on the same day you were supposed to take it	If you miss a dose and do not remember it until the following day
Take the missed dose on that day.	Do not take the missed dose along with the next scheduled dose. Take the missed dose on the next day and extend the number of days in that treatment week.

Example: If you forget to take the Day 3 dose and do not remember it until Day 4, take the Day 3 dose on Day 4, and extend the total number of days in the treatment week by 1 day. If you miss 2 consecutive doses (for example both Day 3 and Day 4 doses), take the missed doses for the next 2 days, and then extend the treatment week by 2 days.

If you have any further questions on the use of this medicine, ask your healthcare professional.

What are possible side effects from using MAVENCLAD?

These are not all the possible side effects you may have when taking MAVENCLAD. If you experience any side effects not listed here, tell your healthcare professional.

Very common side effects: may affect more than 1 in 10 people

- Nausea
- Headache

Common side effects: may affect up to 1 in 10 people

- Cold sores (oral herpes)
- Rash
- Thinning or hair loss
- Fever
- Abdominal pain
- Toothache
- Flu and flu like symptoms
- Cold symptoms
- Bronchitis or other chest infections
- Symptoms of gastroenteritis (diarrhea, vomiting, abdominal pain)
- Back pain
- Anxiety
- Vaginal infection

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Reduction in the number of certain white blood cells, with symptoms such as infections, feeling unusually tired, fever, aches, pain and flu-like symptoms.		✓	
COMMON			
Shingles, with symptoms such as localized 'band' of severe pain and blistering rash, typically on one side of the upper body or the face. Other symptoms may be headache, burning, tingling, numbness or itchiness of the skin in the affected area, feeling generally unwell or fever in the early stages of infection.		✓	
Hypersensitivity (allergic reaction): skin rash, hives, itching, swelling, shortness of breath, fever, wheezing, runny nose, itchy, watery eyes			✓
UNCOMMON			
Liver problems, with symptoms such as nausea, vomiting, stomach pain, loss of appetite, fatigue, dark urine, skin or whites of the eyes turning yellow.		✓	
VERY RARE			
Tuberculosis, with symptoms such as cough that does not go away, fever or loss of weight.		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store MAVENCLAD at room temperature (15°C to 30°C).
- Store this medicine in the original package in order to protect from moisture.
- Keep out of reach and sight of children.

If you want more information about MAVENCLAD:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: <https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>; the manufacturer's website <https://medinfo.emdserono.ca/en>, or by calling 1-888-677-3243.

MAVENCLAD is a trademark of Merck KGaA, Darmstadt, Germany or its affiliates.

This leaflet was prepared by EMD Serono, a Division of EMD Inc., Canada.

Last Revised: October 04, 2021